



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**History of gymnosperms.**—BERRY<sup>13</sup> has published his paper prepared for the symposium on gymnosperms at the meeting of the Botanical Society at the University of California last August. He outlines the paleobotanical evidence in reference to gymnosperms, which extends more or less continuously from the Devonian to the present time. A diagram not only summarizes the data as to the age and relative abundance of the groups, but also indicates the author's suggestions as to phylogeny. The cycadophyte phylum rises directly from the fern stock through Cycadofilicales, which group in turn gives rise to Williamsoniales, Cycadeoidales, and Cycadales. This splitting up of Bennettitales into two phyla seems to be justified by our increasing knowledge of the *Williamsonia* forms. The Cordaitales are represented as having an independent origin from the fern stock and developing the coniferophytes. The araucarians are credited with being the oldest conifer stock, the groups of Pinaceae being left in a more or less problematical position as to age of origin and ancestral group, and among them the Abietineae are regarded as more modern than the Taxodineae and Cupressineae. Taxaceae and the Ginkgoales are left unconnected, the latter beginning in the later Paleozoic, and the former recognized at the base of the Mesozoic.—J. M. C.

**Morphology of Phylloglossum.**—SAMPSON<sup>14</sup> has investigated this much discussed monotypic genus, chiefly with reference to its "annual storage tuber," the so-called "protocorm" of TREUB. As a result of detailed anatomical investigation, the author concludes that this tuber is "a specialized leafy axis, the terminal bud of which functions both as a means of vegetative reproduction and as an organ of perennation," comparable with the resting buds of *Lycopodium inundatum* and the "tubers" of certain Indian species of *Selaginella*. That in fertile plants the tuber is a modified branch is supported by the following facts: (1) a gap is left in the stele of the main axis by the exit of the vascular strand of the tuber; (2) the stele of the tuber often shows a corresponding gap; and (3) the tuber bears leaves, some of which are considerably reduced. The sterile plant consists of a simple axis, the apex of which has formed a storage tuber. The author concludes that the tuber of *Phylloglossum*, therefore, can no longer be compared with the protocorm of *Lycopodium cernuum*, but the two genera are found to be more nearly related by the fact that *Phylloglossum* is shown to be not characteristically an unbranched form.—J. M. C.

**Flavone derivatives in plants.**—SHIBATA and KISHIDA<sup>15</sup> find that aerial parts of high mountain plants are generally much richer in flavone bodies

<sup>13</sup> BERRY, E. W., The geological history of gymnosperms. *Plant World* 19:27-41. figs. 2. 1915.

<sup>14</sup> SAMPSON, K., The morphology of *Phylloglossum Drummondii* Kunze. *Ann. Botany* 30:315-331. figs. 5. 1916.

<sup>15</sup> SHIBATA, K., und KISHIDA, M., Untersuchungen über das Vorkommen und physiologische Bedeutung der Flavorderivate in den Pflanzen. II Mitt. *Bot. Mag. Tokyo* 29:316-332. 1915.